

Electron-Jet-Overlap Study

03/07/05


On the next slide I'm trying to get an idea of how often it happens that a parton is nearby an electron/positron.

Here I'm using the same MC as for the unsmearing studies:

- Z+1j Pythia with 2-to-2 processes, only generator info and particle jets
- Electrons/Positrons are generator level particles with $p_T > 25$, $|\eta| < 1.1$
- Partons are generator level quarks/gluons with $p_T > 20$, $|\eta| < 2.5$

dr_elec_part1


dr_elec_part1
Entries
Mean
RMS



1,226 out of 92,187 = 1.3%

dr_posi_part1

dr_posi_part1
Entries
Mean
RMS



1,224 out of 92,197 = 1.3%

1.3% + 1.3% = **2.6%** of partons are near the electron/positron from Z

03/07/05

Now I'm trying to estimate the acceptance corrections (for each inclusive jet multiplicity) due to the jets/partons that are removed when they are near the electron/positron within $dR=0.4$.




Again, I'm using the same MC as for the unsmearing studies:

- Z+1j Pythia with 2-to-2 processes, only generator info and particle jets
- Electrons/Positrons are generator level particles with $p_T > 25$, $\eta_{\text{et}} < 1.1$
- Partons are generator level quarks/gluons with $p_T > 20$, $\eta_{\text{et}} < 2.5$

To get the correction factors I divide two parton multiplicity histograms:

Numerator = inclusive parton multiplicity for all partons ($p_T > 20$, $\eta_{\text{et}} < 2.5$)

Denominator = incl. parton mult. for all partons outside of $dR=0.4$ w.r.t electron/positron from Z



# partons	correction
1	1.026
2	1.038
3	1.040
4	1.044
5	1.043

03/07/05


In the following I'm trying to see whether a dR cut of 0.4 is too tight:

The next slide looks at the dR distribution between probe-tracks (based on my standard tag-and-probe method) and reco'd CAL jets (passing all the usual jet quality cuts):


- top row is Pythia MC
- bottom row is data
- left column is w/o elec-jet-overlap cut
- right column is with elec-jet-overlap cut ($dR=0.4$)

tag-and-probe method: dR between probe-track and all good jets


dr(elec-CALjet): w/o jet-elec overlap cut




dr(elec-CALjet): with jet-elec overlap cut



dr(elec-CALjet): w/o jet-elec overlap cut



dr(elec-CALjet): with jet-elec overlap cut



On the next slide I'm trying to understand how often it happens that a parton that is **outside** the $dR=0.4$ elec-jet-overlap-cone, has a reco'd CAL jet **inside** the cone, i.e. the jet algorithm merges the fake jet from the electron with the real jet so that the final jet is 'pulled' inside the cone.

This is done using regular Pythia MC.



Electrons/Positrons are generator level particles with $p_T > 25$, $\text{le}^{\text{rel}} < 1.1$

Partons are generator level quarks/gluons with $p_T > 20$, $\text{le}^{\text{rel}} < 2.5$

CALjets are reco'd CAL jets passing all quality cuts.

Numerator = $dR(\text{generated electron/positron} - \text{parton})$

Denominator = $dR(\text{generated electron/positron} - \text{CALjet})$



Looks like a large number of jets between $dR=0.5$ and $dR=0.6$ get 'pulled' away!

03/07/05